

Assessment of the nutritional knowledge and nutrient Intake of lactating women and its impact on body mass index

**Ahmad, E. Shoman*, Amany, M .S. Abd Elhay*,
Hany, H.A. Elgazzar** and Amira, S. Nassar****

*Faculty of Medicine, Ain Shams University, ** National Nutrition Institute

Abstract

The study was done to assess the nutritional knowledge and nutrients intake among lactating women. The total sample size was 151 lactating women selected randomly from those attending the Breastfeeding clinic at the center for social and preventive medicine (CSPM). The results revealed that obesity was the dominant feature in 47.7% of cases, while overweight and normal weight constituted 26.5% and 25.8% respectively. The majority of the women had very good nutrition knowledge (77.5%), while 22.5% had a good knowledge. 76.2% women were at risk of hunger and 23.8 were food secure. Poor maternal dietary quality may have implications for both mother and child, and socioeconomic position and nutritional knowledge are likely to be important. Effective interventions to support mothers to achieve healthy diets for themselves and their families are needed. Lactation period is an opportunity to increase awareness of the mother toward healthy nutritional practices. Achievement and maintenance of good dietary habits is expected to duplicate the benefit of health outcomes when mothers become models for their children and their families. Nutrition education for mothers should be provided in the all health and PHC centers that deal with breastfeeding mothers. It also important to increase awareness of lactating mother's and educate them about the quality of diet through mass media.

Introduction

Lactation is the medical term for yielding milk by the mammary glands to practice breastfeeding. (*Wosje et al, 2004*). Exclusive breastfeeding means that the infant receives only breast milk, no other liquid or solids are given not even water with exception of oral rehydration solution or drops of vitamins, minerals or medicine (*WHO, 2013*).

Exclusive breastfeeding is recommended by all the international agencies for the first six months of life. Hence, the nutritional status of the lactating mother is considered to be important for the well being of both mother and infant (*Allen, 2005*). Because of the immature intestinal immune function, the newborn is susceptible to intestinal and systemic infection; exclusively breastfed infants are less likely to develop infection than bottle fed infants (*Walker, 2010*). It protects against diarrhea, pneumonia, and have a longer term benefit; as decrease mean blood pressure, cholesterol, decrease the prevalence of obesity and type 2 diabetes (*US department of health and human services, 2005*).

It is shown that 15% of the global burden disease is attributed to the combined effects of child and maternal underweight or micronutrient deficiencies. Women at reproductive age are vulnerable to malnutrition since they have special nutrients needs (*WHO, 2008*).

The levels of some nutrients such as thiamine, riboflavin, vitaminB-6, vitaminB-12, iodine, and selenium in breast milk are reflection of the dietary intake of mothers. Other micronutrients (such as folic acid, calcium, iron, copper, and zinc) remain at relatively high levels in breast milk even when the mother's reserves are low (*Allen, 2005*).

Carbohydrates are essential to provide the lactating mother with energy during the nursing period. Carbohydrates are also necessary in the diet to spare the utilization of body protein and prevent ketosis. Lactating mother needs about 80g/day carbohydrates over what is recommended for pregnancy (*Thompson and Mannore, 2005*).

Protein is responsible for various function, i.e. cell growth, tissue repair, a strong immune system, and may be a source of energy if needed. Lactating mothers need 10 to 15 g/day protein over the needs of non-pregnant women (*labbok et al,1990*).

Monounsaturated fats are important, and polyunsaturated fats that practically contains omega-3 fatty acids. It is important for the infant's central nervous system development and also brain growth and eye development (*Lawrence , 2008*). Therefore, mothers should take a balanced daily diet to replenish their stores(*US department of health and human services, 2005*).

Mothers are the major providers of food for their families and are also a substantial source of nutrition information for their children. It is thus important that mothers should have good nutrition knowledge and be aware of the recommended intakes of core food (*WHO, 2008*). Nutrition knowledge is the understanding of different types of food and how food nourishes the body and influences health .Although breastfeeding is one of the most natural functions of a women's body, knowledge about lactation can make breastfeeding a success for both the mother and infant (*Insel , et aL,2003*).

In developing countries, the prevalence of several micronutrients deficiencies may cause poor maternal infant interaction and affect the mental status of the mother during breastfeeding (*Frith and Naved, 2009*). The nutritional status of women in these countries is usually threatened by repeated un-spaced pregnancies which may lead to several nutrient deficiencies. Egypt is one of the developing countries; food consumption studies show that the intake of vitamin A, calcium, iron, iodine, selenium, and zinc are insufficient. Cereals represent the main source of energy in Egypt diet (*FAO, 2011*).

Therefore it is justified to call for nutrition interventions for further improvement of health of lactating mothers and their infants (*Haidar,et al.,2003*).

Aim of the study:

The study was designed to show the practice of the lactating mothers concerning breastfeeding for determining any current possible variation and to give recommendations to them for nutrition support for the mother and child.

Subjects and Methods

Study design: Descriptive cross-sectional study.

Study site: Breastfeeding clinic at the center for social and preventive medicine (CSPM) of Abu El Reish children's Hospital.

Study time: from July 2014 till January 2015

Study population:

A group of lactating women who accepted to participate in the study were interviewed. Two participants were selected randomly, on every working day, from the lactating women who fulfilled the inclusion criteria and who accepted to participate.

Inclusion Criteria: The women were considered lactating if they breastfed their infant on demand.

Sample size: 151 lactating women were included in the study.

Study tools:

I- Interview questionnaires: The questionnaires, were adapted from standardized questionnaires of previous study (Nyaboke A, 2010), to match with our Egyptian culture. The quantitative food frequency questionnaire (QFFQ) was adapted from a validated food frequency questionnaire developed by the National Nutrition Institute.

Five types of questionnaires have been used;

1- Socio-bio Demographic Questionnaire (S-BDQ); this provided information on factors relevant to lactating woman regarding the environment she lived in. The data collected included age of the mothers, occupation, marital status, educational level, and number of children as well as number of people in the household. Data on socio-economic status were gathered about housing standards, food, water supply, and fuel sources, daily family expenditure on food and average monthly income for family while information on health status was covered by the number visits to out-patient and in-patient clinics.

2- Nutritional knowledge; the questionnaire covered the knowledge of the mothers about the varies nutrients in food groups. A score of four or less (0-4) indicated poor nutrition knowledge. A score of five to eight indicated that the mother had good nutrition knowledge. A score of nine to 12 indicated that the mother had very good nutrition knowledge.

3- The Quantitative Food Frequency Questionnaire (QFFQ); this provided information on the eating pattern and intake. The Food Frequency questionnaire comprises a list of foods and beverages on which respondents, reported their usual amount and frequency of consumption (WHO, 2014).

4- The Hunger Scale; provided information on the food availability and the consumption by the lactating mothers and their households. The Hunger Scale questionnaire has three main components that might be used to determine domestic hunger. These are; household level insecurity, individual level insecurity and child hunger. It consists of eight questions, a score of five positive (yes) or more indicates a food shortage problem affecting everyone in the household. These families were considered as "hungry". A score of one to four indicates that the family is at risk of hunger (Labadarios, et al., 1999).

5- 24-hr Recall Questionnaire (24-H-RQ); this provides information on the types of foods consumed at mealtimes and between meals, portion sizes over the past twenty four hours. This information is then used to calculate mean daily supplies of nutrient.

II- Anthropometric Measures: weight and height were measured and then BMI was calculated for each participant.

Weight measurement procedures for participant were in accordance to (Hammond, 1998) as follows:

- A calibrated level plate from scale placed on a flat hard surface was used.
- Participants were measured without shoes.

- Participants were directed to stand in the center of the platform, with the body weight equally distributed between both feet.

Height measurements Procedures were performed according to (*Hammond, 1998*).

-Height was measured while the participants were without shoes
-The participant was directed to stand on a flat surface with weight distributed evenly on both feet, heels together and head positioned so that the line of vision was perpendicular to the body.

-The arms hang freely and the head, back, buttock, and heels were in contact with vertical board.
-An attached metric ruler and a movable horizontal head board was used.
-After the participant stood properly, the head board was moved down to the top of head so that the hair was compressed.

Body mass index (BMI) was calculated using the following formula: **Weight (Kgs)/height (m) ²**.

The BMI of the participants was classified according to WHO, 1995 as;

- 1) Underweight if BMI is less than 18.5 Kg\m²
- 2) Normal if BMI is between 18.5- 24.9 Kg\m²
- 3) Overweight if BMI is between 25-29.9kg\m².
- 4) Obese if BMI is 30.0kg\m².

Results

Table (1) shows the socio-demographic characters of the lactating mothers. The mean age of the lactating women was (28.6), majority of the women were married (98%) and those who received high education were (43.6%). The majority of the mothers were housewives (51.7%) followed by the employees (31.2%) then the professional (11.9%).

Distribution of the mothers according to the monthly income and the money spent on food per week table (2), revealed that 52.3% of the families earned from 1001-3000 L.E, while 49.7% had spent from 100-200-L.E on food per week, and those who used to spend more than 500L.E were (7.3%).

Table (3) illustrates the distribution of the mothers according to the nutritional knowledge score and hunger score of the lactating women. It was clear that 77.5% had a very good knowledge score and 22.5% had a good nutritional knowledge score. However, 76.2% of the lactating women were at risk of hunger and 23.8% were food secure.

Table (4) illustrates the mean 24hr recalls intake among the lactating women; it revealed that the consumption of carbohydrates and protein covered the requirements while there was over consumption of fat. Most the micronutrients was below the recommendations as vitamin A, vitamin B12, vitamin B6, vitamin B2 and calcium.

Distribution of the lactating mothers by dietary adequacy of some nutrients as shown in table (5). The majority of the women (89.4%) consumed >100% which is over consumption, and only 2% consumed <50% of the recommended daily allowance (RDA) from carbohydrates. Over half of the women 55.1% consumed .100% of the RDA Of protein, 34% consumed 75-100% which is acceptable, and 10.9% between 50-75% of the recommended protein intake.

Table (6) revealed that (17.7%) of the study group take their acceptable RDAs from calcium while (62.6%) had taken unacceptable (50%-75%) or unsafe levels (<50%) of their requirements. Regarding iron (91.4%) had overconsumption and only (2.6%) had less than 50%. (43.0%) of the study group consumed their acceptable RDAs from Zinc. Seventy nine Of the study group consumed Vit.B12 below 50% of the recommended daily allowances (RDAs) and (75) of the study group consumed iodine below 50% of the RDAs table (7).

Table (8) shows the results of the BMI of the lactating women, we noticed that obesity was more prominent 47.7% , while normal weight and overweight was 25.8% and 26.5% respectively.

Discussion

A study was done by **Shalabi (2003)** who assessed the nutrition status of lactating women in some rural areas in Menufyia governorate and found that the percent of the lactating mothers in the age group < 25 years represented 45.5%. In addition the highest percent of the lactating mothers were housewives. Additionally, finding of this study coincide with those of **Heikel (2010)** who conducted a study at the same CSPM and found that 17% of mothers were illiterate and 41% had a secondary or higher education. Furthermore, 90% of the studied participants were not working for cash (housewives).

The Egyptian Demographic and Health Survey (E-DHS), 2008 reported similar findings to ours where 60.5% of Egyptian families have less than five individuals and the average household had 4.6 individuals per household (**El Zanaty and Way, 2009**)

The present study revealed that the majority of the participant's family had earned from 1001-3000 L.E monthly income, and 49.7% of the families have spent 100-200L.E on food per week, (table:1-3). A recent review of the food situation in Egypt by the World food programme (2011) concluded that the key underlying and basic factors of household food insecurity in Egypt include; low income to allow access to more diversified foods including fruits, vegetables and animal protein leading to unbalanced diets which are often high on energy but low in micro-nutrients, Increased cost of living and Sub-optimal access to social assistance systems including the ration card for subsidized staple foods and cash assistance.

From our study, the majority of the mothers had a very good nutritional knowledge and or a good knowledge and this could be explained that a good number of the participants of the study group had a high university education and the rest might had the knowledge from the clinic or the T.V. A study was done by **Mcloed, 2011** in England reported that Overall nutrition knowledge was found to be quite high, with mean nutrition knowledge scores of 12.5, 12.7, and 13.7 out of a possible 17, in the low, medium, and high socioeconomic groups, respectively. Maternal nutrition knowledge was found to partly mediate the association between socioeconomic position and maternal diet quality.

The present study agrees with a study done by **Asfaw (2007)** who investigated the relationship between micronutrient deficiency and the prevalence of mothers' overweight/obesity in Egypt using the 1997 Egyptian Integrated Household Survey. This study published that the prevalence of overweight/obese is 80.8% higher for micronutrient deficient mothers than for non-deficient mothers, keeping all other variables constant. Egyptian food subsidy program, which lowers the relative prices of energy-dense, nutrient-poor food items, can be one of the major factors for the emergence of overweight/obese and micronutrient deficient mothers in the country.

Conclusion

Lactation period is an opportunity to increase awareness of the mother toward healthy nutritional practices for the benefit of the mothers and child health. Effective interventions to support mothers to achieve healthy diets for themselves and their families are needed.

**Table (1):
The sociodemographic characteristics of the study group.**

Lactating mothers	Mean \pm SD	Min - Max
Age in years	28.6 \pm 5.5	18.0 – 38.0
Baby age in months	8.1 \pm 3.9	2.0 – 18.0
	No	%
Marital status		
Married	148	98.0
Divorced	3	2.0
Education level		
Illiterate	19	12.6
Primary	11	7.3
Preparatory	14	9.3
Secondary	41	27.2
University	66	43.6
Occupation		
Housewives	78	51.7
Non-skilled	4	2.6
Skilled	4	2.6
Employees	47	31.2
Professional	18	11.9
No of the family members		
3	33	21.9
4	46	30.5
5	54	35.7
6	16	10.6
> 6	2	1.3
Total	151	100%

Table (2):
Distribution of the mothers according to the monthly income,
the money spent on food per week.

Income and expenditure	No	%
Monthly income (L.E)		
< 500	6	4.0
500-1000	48	31.8
1001-3000	79	52.3
3001-5000	6	4.0
> 5001	12	7.9
Money spent on food per week(L.E)		
< 100	22	14.6
100-200	75	49.7
201-300	36	23.8
301-500	7	4.6
> 500	11	7.3

Table (3):
Distribution of the mothers according to the nutritional knowledge score
and hunger scale score of the study group.

Parameters	No	%
Nutritional knowledge score		
Very good	117	77.5
Good	34	22.5
Poor	0	0
Hunger scale score		
At risk of hunger	115	76.2
Food secure	36	23.8
Hunger	0	0

Table(4):
Mean of 24 hours intake among the study group.

Intake	Mean \pm SD	WHO recommendations
Calories	2900 \pm 440	2803
Protein(gm.)	74.3 \pm 18.3	71
Carbohydrate(gm.)	301.0 \pm 69.4	210
Fat(gm.)	65.8 \pm 23.0	27
Calcium(mg.)	721 \pm 343.6	1300
Iron(mg.)	20 \pm 7.6	10
Magnesium(mg)	146 \pm 62.0	270
Vitamin A(μ g)	560 \pm 733.3	1300
Vitamin D(μ g)	4.11 \pm 8.7	5
Vitamin C(mg)	169 \pm 150	120
B 1(thiamine)(mg)	1.1 \pm 0.5	1.4
B 2(riboflavin)(mg)	0.96 \pm 0.6	1.6
B 6(mg)	0.14 \pm 0.1	2
B 12 (μ g)	0.29 \pm 0.4	2.8

Table (5):
Percentage distribution of the lactating mothers by dietary adequacy of some nutrients

	No	%
<u>Carbohydrate</u>		
Unsafe (< 50%)	3	2.0
Unacceptable (50 – 75%)	6	4.0
Acceptable (75 – 100%)	7	4.6
Overconsumption (> 100%)	135	89.4
<u>Calories</u>		
Unsafe (< 50%)	11	7.3
Unacceptable (50 – 75%)	54	35.7
Acceptable (75 – 100%)	77	51.0
Overconsumption (> 100)	9	6.0
<u>Protein</u>		
Unacceptable (50 – 75%)	19	10.9
Acceptable (75 – 100%)	51	34.0
Overconsumption (> 100)	81	55.1

Table (6):
Percentage distribution of the lactating mothers by dietary adequacy of some nutrients

	No	%
<u>Calcium</u>		
< Unsafe (< 50%)	41	27.9
Unacceptable (50 – 75%)	51	34.7
Acceptable (75 – 100%)	26	17.7
Overconsumption (> 100)	29	19.7
<u>Iron</u>		
< Unsafe (< 50%)	4	2.6
Unacceptable (50 – 75%)	6	4.0
Acceptable (75 – 100%)	3	2.0
Overconsumption (> 100)	138	91.4
<u>Zinc</u>		
< Unsafe (< 50%)	14	9.3
Unacceptable (50 – 75%)	29	19.2
Acceptable (75 – 100%)	65	43.0
Overconsumption (> 100)	43	28.5
<u>Selenium</u>		
Unsafe (< 50%)	75	100.0

Table (7):
Percentage distribution of the lactating mothers by dietary adequacy of some nutrients

	No	%
<u>Vit.B12</u> < Unsafe (< 50%)	79	100.0
<u>Vit.A</u> < Unsafe (< 50%)	115	76.2
Unacceptable (50 – 75%)	19	12.6
Acceptable (75 – 100%)	10	6.6
Overconsumption (> 100)	7	4.6
Iodine < 50%	75	100.0
<u>Vit.D</u> < 50%	121	80.1
> 100%	30	19.9
<u>Vit.C</u> < 50%	147	97.4
50 – 75%	4	2.6

Table (8):
The anthropometric measurements of the study group

	Mean ±SD	Min - Max
Weight	75.1±14.0	45.0 – 107.0
Height	160.0 ±5.6	145.0 – 174.0
MBI	29.3±5.2	18.6 – 42.9
	No	%
BMI categories		
Normal weight(18.5-24.9kg/m ²)	39	25.8
Overweight (25.0-29.9kg/m ²)	40	26.5
Obese(30kg/m ²)	72	47.7

References

Allen, LH, (2005):

Multiple micronutrients in pregnancy and lactation: an overview. *American Journal of Clinical Nutrition*, 81(Supply): 1206S

Asfaw A (2007):

Micronutrient deficiency and the prevalence of mothers' overweight/obesity in Egypt¹ International Food Policy Research Institute (IFPRI), Washington, DC 20006, USA

El-zanaty F and Way A (2009):

Egypt Demographic and Health Survey 2008. Cairo, Egypt: Ministry of Health, El-Zanaty and Associates and Macrointernational

(FAO) Food and Agricultural Organization, (2011):

The double burden of malnutrition. Case studies for six developing countries. Policies and strategies to improve nutritional status. URL:<http://www.fao.org/docrep/009/a0442e/a0442e00.htm> Retrieved November 13th.

Frith AL, Naved RT, (2009):

Micronutrient supplementation affects maternal infant feeding interaction and maternal distress in Bangladesh. *Am J Clin Nutr*, 90(1):141 – 8.

Haidar J, Muroki NM, Omwega AM, Ayana G, (2003):

Malnutrition and iron deficiency in lactating women in urban slum communities from Addis Ababa, Ethiopia. *East African Medical Journal*, 80(4):191 – 194.

Hammond K., (1998):

The history and physical examination for nutritional assessment. *Public Health Nut*, 5: 366 – 375.

Heikel S (2010):

Assessment of quality of services provided by antenatal care clinic in Center of Social and Preventive Medicine. MSc thesis in Public Health, Cairo University

Insel P, Turner E and Ross D (2003):

Discovering nutrition: Jones and Bartlett (eds) USA Thompson Wadsworth.

Labbok M, Mcdonald M, Besely M, Klyenkya, Isabiye M, Greavevs-J, O'Garac,(1990):

Proceedings of the Interagency workshop on health care practices related to Breastfeeding. *International Journal of Gynecology and Obstetrics* ;31(1).

Labadarios D., stayon N., Emawnder, Maclytyre U, Swart R, Gortcke,Huskisson A ., Haosten,(1999):

National Food Consumption survey of aged 1.9 years in south Africa.

Lawerence R., (2008):

Encyclopedia and curly childhood Development: Breastfeeding Colorado's; 2008.

McLeod ER, Campbell KJ and Hesketh KD (2011):

Nutrition Knowledge; a mediator between socioeconomic position and diet quality in Australian first-time mothers; 111(5):696-704. doi:10.1016/j.jada.02.011.

Nyaboke A., (2010):

Nutrient Intake and Nutrition Knowledge of Lactating Women (0-6 months postpartum) in a Low Socio – Economic Area in Nairobi, Kenya. Thesis submitted in partial fulfillment for the requirement of Master of Science Degree In Human Nutrition in the Faculty of Natural and Agricultural Sciences.

Shalabi RA (2003):

Nutritional assessment of lactating mothers in some rural areas in Menufyia. MSc thesis in Nutrition and Food Science. Home Economics, Menufyia University

Thompson J and Mannore M, (2005):

Nutrition ,An applied approach. Newyork, Pearson Corporation; 3.1(2):55.

U.S. Department of Health and Human Services and U.S. Department of Agriculture, (2005):

Dietary Guidelines for Americans, 2005. 6th Edition, Washington, DC: U.S. Government Printing Office. URL: <http://www.health.gov/dietaryguidelines/dga2005/document/pdf/DGA2005.pdf> Retrieved 9th June, 2011.

Walker A, (2010):

Breast milk as the gold standard for protective nutrients. The Journal of Pediatrics, 156 (2), Supplement (1): S3 – S7.

World and Health Organization(WHO),(1995):

Physical status. The use and interpretation of anthropometry. Report of a WHO expert committee. WHO technical report series number 854. Geneva: World Health organization. URL: http://www.who.int/childgrowth/publications/physical_status/en/index.html Retrieved 3rd February, 2012.

World and Health Organization (WHO), (2008):

10 Facts about women's health: Geneva: available from: <http://www.who.org/women;health>. Accessed on 4th June .

World and Health Organization (WHO),(2013):

Exclusive breast feeding : http://www.who.int/elena/titles/exclusive_breastfeeding/en/ 23 May 10:53 CEST

World and Health Organization (WHO),(2014) :

WHO guidelines on nutrition http://www.who.int/publications/guidelines/reproductive_health/en/. Accessed on June 21.

Wosje K S and Kalkwarf H J ,(2004):

Lactation, weaning, and calcium Supplementation: effects on body composition in postpartum women. American Society for clinical Nutrition August 80: 2 423-429

تقييم الوعي التغذوي والتناول الغذائي للأمهات المرضعات
و تأثيرهما على مؤشر كتلة الجسم

د. احمد عصمت شومان*، د. امانى السيد*، د. هانى الجزار**، د. اميرة سعيد**

*كلية الطب، جامعة عين شمس، **المعهد القومى للتغذية

الملخص العربى

تم إجراء هذه الدراسة لتقييم المعرفة الغذائية و العناصر المغذية بين النساء المرضعات. بلغ مجموع حجم العينة ١٥١ تم اختيارها عشوائيا من أولئك الذين يحضرون عيادة الرضاعة الطبيعية في مركز للطب الاجتماعي والوقائي . و كشفت النتائج أن البدانة كانت السمة الغالبة في ٤٧,٧% من الحالات، في حين يشكل زيادة الوزن و الوزن الطبيعي ٢٦,٥% و ٢٥,٨% على التوالي. فالنسبة للمعرفة الغذائية كانت ٧٧% جيدة جدا و ٢٢,٥% كانت المعرفة الغذائية جيدة. كانت ٧٦,٢% من النساء المرضعات المعرضين لخطر الجوع وكانت ٢٣,٨% منهن مكتفين غذائين. فترة الرضاعة هي فرصة لزيادة وعي للأم تجاه الممارسات الغذائية السليمة. و من المتوقع أن يتضاعف فائدة الوعي الصحى عندما تصبح الأمهات نماذج لأطفالهم وأسرهـم لتحقيق وصول العادات الغذائية الجيدة. و ينبغي توفير التثقيف الغذائي للأمهات في جميع المراكز الصحية والرعاية الصحية الأولية التي تتعامل مع الأمهات المرضعات و من المهم أيضا لزيادة الوعي الأم المرضعة و تثقيفهم حول نوعية النظام الغذائي من خلال وسائل الإعلام.